


AD 723515

ACIC - TC - 1679

# TECHNICAL TRANSLATION

THE PROBLEM OF REDUCING THE FORMULA DESCRIBING THE SHAPE OF THE  
EARTH TO A TAYLOR SERIES


By: W.I. Marych



Distribution of this document  
is unlimited.

AERONAUTICAL CHART AND INFORMATION CENTER  
RESEARCH DIVISION LIBRARY BRANCH  
TECHNICAL LIBRARY SECTION  
ST. LOUIS, MISSOURI 63118

Reproduced by  
**NATIONAL TECHNICAL  
INFORMATION SERVICE**  
Springfield, Va. 22151



Unclassified

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author, ...)		2a. REPORT SECURITY CLASSIFICATION	
Aeronautical Chart & Information Center		Unclassified	
		2b. GROUP	
3. REPORT TITLE			
THE PROBLEM OF REDUCING THE FORMULA DESCRIBING THE SHAPE OF THE EARTH TO A TAYLOR SERIES.			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Source: Geodeziya, Kartografiya i Aerofotos"yemka, Vol. 9, 1969, pp. 29-32, Mezvedomstvennyy Respublikanskiy Nauchno-tehnicheskii			
5. AUTHOR(S) (First name, middle initial, last name) Sbornik, Lvov, Ukrainian SSR.			
Marych, M. I.			
6. REPORT DATE		7a. TOTAL NO. OF PAGES	7b. NO. OF REFS
November 1970		6	2
8a. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER(S)	
b. PROJECT NO.		ACIC-TC-1679	
c.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.			
10. DISTRIBUTION STATEMENT			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY	
		This document has been approved for public release and sale; its distribution is unlimited.	
13. ABSTRACT			
<p>Molodenskiy has suggested a new method of solving the integral equation for the density of a simple layer and has obtained formulae for successive approximations of the disturbing potential on the physical surface of the earth. If the earth's physical surface on which the gravity anomalies are given is assumed to be a sphere, then the disturbing potential will be determined by Stokes' formula. This paper shows an attempt to obtain this result using the Molodenskiy method of successive approximations, and also the method of transformations by expanding the disturbing potential in a Taylor series.</p>			

DD FORM 1473  
1 NOV 65

Unclassified

Security Classification



TRANSLATION

ACIC-TC- 1670

THE PROBLEM OF REDUCING THE FORMULA DESCRIBING THE SHAPE OF THE EARTH  
TO A TAYLOR SERIES

(K voprosu privedeniya formuly, opredelyayushchey figuru Zemli, k ryadu  
Teylora)

By: Marych, M.I.

Source: Geodeziya, Kartografiya i Aerofotos"yemka, Vol. 2, 1960, pp.  
29-32, Mezhdomeystvennyy Respublikanskiy Nauchno-tehnicheskiy  
Sbornik, Lvov, Ukrainian SSR.

Translated by: George Synenko, Technical Translation Unit, November, 1970

Distribution of this document  
is unlimited.

TECHNICAL LIBRARY SECTION  
LIBRARY BRANCH  
RESEARCH DIVISION  
AERONAUTICAL CHART AND INFORMATION CENTER  
ST. LOUIS, MISSOURI 63118

# NOTICE

THIS TRANSLATION IS A RENDITION OF THE ORIGINAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE AERONAUTICAL CHART AND INFORMATION CENTER.

TC DENOTES TRANSLATION CODE SYSTEM UTILIZED BY AERONAUTICAL CHART AND INFORMATION CENTER. WHEN ORDERING A TRANSLATION REFER TO TC NUMBER.

THE PROBLEM OF REDUCING THE FORMULA DESCRIBING THE SHAPE  
OF THE EARTH TO A TAYLOR SERIES

M.S. Molodenskiy has suggested a new method of solving his integral equation for the density of a simple layer and has obtained formulae for successive approximations of the disturbing potential on the physical surface of the earth [2]. Two first approximations, as is shown in work [1], are reduced to the sum of the first two terms of a Taylor-series expansion of the disturbing potential in powers of the earth relief height, where the vertical gradients of the gravity anomalies are calculated according to Numerov's formula. If we assume the earth's physical surface on which the gravity anomalies are given to be a sphere, then the disturbing potential, as is known, will be determined by Stokes' formula. Let us try to obtain this result, using the Molodenskiy method of successive approximations, and also the method of transformations by expanding the disturbing potential in a Taylor series.

1. Let us find by Molodenskiy's method the values of the disturbing potential on the sphere of radius  $\varrho = a + H$ , where  $a$  is the mean radius of the earth. First of all let us note that with  $H = \text{const}$  Molodenskiy's formula coincides with Stokes' formula. However, the integration here is carried out not with respect to the sphere of radius  $\varrho$ , but with respect to the sphere of radius  $a$ . This is explained by the fact that in deriving the formula for the general case, small quantities depending on  $H$  were discarded. Consequently, let us proceed from the integral equation

$$2\pi\varphi = \Delta g + \frac{3}{2} \int \varphi \frac{d\sigma}{r},$$

where  $\Delta g$  = gravity anomaly;

$$r = 2 \sin \frac{\psi}{2};$$

$\psi$  = angle formed by the radius vectors of the given and current points;  
 $d\sigma$  = surface element of the unit-radius sphere, density of simple layer distributed on the sphere of radius  $\varrho$ .

Introducing the new function

$$\chi = \frac{\rho^3}{a^2} \varphi,$$

we have

$$2\pi\chi = \frac{\rho^3}{a^2} \Delta g + \frac{3}{2} \int \chi \frac{d\sigma}{r}.$$

By means of the small parameter  $k(0 \leq k \leq 1)$  we may write

$$2\pi \sum_{n=0}^{\infty} k^n \chi_n = \frac{\bar{\rho}^2}{a^2} \Delta g + \frac{3}{2} \int \sum_{n=0}^{\infty} k^n \chi_n \frac{dz}{r},$$

where  $\bar{\rho} = a + kH$ . The series entering into the left and right parts of the equation must be equal for all values of  $k$ ; therefore the multipliers with  $k^n$  in both parts of the equation must be mutually equal. Thus, we obtain

$$2\pi \chi_n - \frac{3}{2} \int \chi_n \frac{dz}{r} = G_n,$$

where

$$G_0 = \Delta g, G_1 = \frac{2H}{a} \Delta g, G_2 = \frac{H^2}{a^2} \Delta g, G_{n>3} = 0.$$

Solving these integral equations, we find

$$\chi_n = \frac{G_n}{2\pi} + \frac{3}{(4\pi)^2} \int G_n s(\psi) dz,$$

and also

$$\int \chi_n \frac{dz}{r} = \frac{1}{4\pi} \int G_n s(\psi) dz, \quad (1)$$

where  $s(\psi)$  is the Stokes function. The disturbing potential determined by the formula

$$T = \frac{a^2}{\rho} \int \chi \frac{dz}{r}, \quad (2)$$

is found, as well as the subsidiary function  $\chi$ , by successive approximations, i.e.,

$$T = \sum_{n=0}^{\infty} T_n,$$

where each term  $T_n$  contains the height  $H$  only to the degree  $n$ . Thus, taking into consideration expressions (1) and (2) we obtain

$$\begin{aligned}
T_0 &= a \int \chi_0 \frac{d\sigma}{r} = \frac{a}{4\pi} \int \Delta g s(\psi) d\sigma; \\
T_1 &= a \int \chi_1 \frac{d\sigma}{r} - H \int \chi_0 \frac{d\sigma}{r} = \frac{H}{4\pi} \int \Delta g s(\psi) d\sigma; \\
T_2 &= a \int \chi_2 \frac{d\sigma}{r} - H \int \chi_1 \frac{d\sigma}{r} + \frac{H^2}{a} \int \chi_0 \frac{d\sigma}{r} = 0; \\
T_{n>3} &= 0.
\end{aligned}$$

Consequently:

$$T = \frac{\rho}{4\pi} \int \Delta g s(\psi) d\sigma. \quad (3)$$

Thus, if the anomalies  $\Delta g$  are given on the sphere, then Molodenskiy's process of successive approximations leads to Stokes' formula.

2. Let us use the expansion of the disturbing potential in a Taylor series

$$T = \frac{a}{4\pi} \int \left( \Delta g - H \frac{d\Delta g}{d\rho} + \frac{H^2}{2} \frac{d^2 \Delta g}{d\rho^2} - \dots \right) s(\psi) d\sigma + \frac{dT}{d\rho} H - \frac{1}{2} \frac{d^2 T}{d\rho^2} H^2 + \dots \quad (4)$$

and find the values  $T$  at points on the sphere of radius  $\rho$ . First let us obtain auxiliary relations. For this purpose we take into consideration Molodenskiy's integral formula

$$\frac{dU_0}{d\rho} = \frac{1}{2\pi\rho} \int \frac{U - U_0}{r^3} d\sigma + \frac{U_0}{\rho},$$

which determines the value of the normal derivative for the harmonic function  $U$ , given on the surface of the sphere [2]. Applying this formula to the harmonic functions

$$\rho \frac{dT}{d\rho}, \rho \frac{d}{d\rho} \left( \rho \frac{dT}{d\rho} \right), \dots,$$

where

$$\frac{dT}{d\rho} = -\Delta g - \frac{2T}{\rho}, \quad (5)$$



we find

$$\frac{d^i \Delta g}{d\rho^i} = \frac{1}{2\pi\rho} \int \left[ \frac{d^{i-1} \Delta g}{d\rho^{i-1}} - \left( \frac{d^{i-1} \Delta g}{d\rho^{i-1}} \right)_0 \right] \frac{d\sigma}{r^3} - (i+1) \frac{d^{i-1} \Delta g}{\rho d\rho^{i-1}}. \quad (6)$$

(i = 1, 2, ...)

Presenting the quantity  $\frac{d^{i-1} \Delta g}{d\rho^{i-1}}$  in the form of a series expansion in spherical harmonics and considering that

$$r^{-3} = -\frac{1}{2} \sum_{n=0}^{\infty} (2n+1)(n+1) P_n(\psi),$$

where  $P_n$  are Legendre's polynomials of the nth order, after corresponding transformations we obtain

$$\frac{d^i \Delta g}{d\rho^i} = -\frac{1}{\rho} \sum_{n=0}^{\infty} (n+i+1) \left( \frac{d^{i-1} \Delta g}{d\rho^{i-1}} \right)_n. \quad (7)$$

These relations, and also

$$s(\psi) = \sum_{n=2}^{\infty} \frac{2n+1}{n-1} P_n(\psi) \quad (8)$$

make it possible without particular difficulties to carry out the integration indicated in formula (4). Since

$$\frac{a}{4\pi} \int \frac{d\Delta g}{d\rho} s(\psi) d\sigma = \frac{\rho}{4\pi} \int \frac{d\Delta g}{d\rho} s(\psi) d\sigma - \frac{H}{4\pi} \int \frac{d\Delta g}{d\rho} s(\psi) d\sigma,$$

then with the help of (7) and (8) we find

$$\frac{a}{4\pi} \int \frac{d\Delta g}{d\rho} s(\psi) d\sigma = -\Delta g - \frac{3T}{\rho} + \frac{H}{\rho} \left( \Delta g + \frac{3T}{\rho} \right), \quad (9)$$

where  $T$  is determined by Stokes' formula (3). In the same way we find

$$\frac{a}{4\pi} \int \frac{d^2 \Delta g}{d\rho^2} s(\psi) d\sigma = -\frac{d\Delta g}{d\rho} + \frac{4\Delta g}{\rho} + \frac{12T}{\rho^2} + \frac{H}{\rho} \left( \frac{d\Delta g}{d\rho} - \frac{4\Delta g}{\rho} - \frac{12T}{\rho^2} \right). \quad (10)$$

Differentiating function (3) in the direction of  $\rho$ , we have

$$\frac{d^2 T}{d\rho^2} = -\frac{d\Delta g}{d\rho} + \frac{2\Delta g}{\rho} + \frac{6T}{\rho^2}. \quad (11)$$

Let us now perform the calculations according to formula (4). With the help of expressions (5) and (9) we obtain

$$\frac{a}{4\pi} \int \left( \Delta g - H \frac{d\Delta g}{d\rho} \right) s(\psi) d\psi + \frac{dT}{d\rho} H = T + \delta_2,$$

where

$$\delta_2 = -\frac{H^2}{\rho} \left( \Delta g + \frac{3T}{\rho} \right).$$

The given result differs from the exact value  $T$  of the disturbing potential by the quantity  $\delta_2$  containing  $H^2$ . If in formula (6) with  $i = 1$  it is assumed  $\rho = a$ , which corresponds to the calculation of the vertical gradient of the gravity anomaly according to Numerov's formula, then instead of formula (9) we obtain

$$\frac{a}{4\pi} \int \frac{d\Delta g}{d\rho} s(\psi) d\psi = -\Delta g - \frac{3T}{\rho},$$

and consequently the quantity  $\delta_2$  will not appear in the result. This fully agrees with the condition that the two first Molodenskiy approximations consider quantities which contain  $H$  only to the first degree. Further, taking into consideration (10) and (11) we have

$$\frac{a}{4\pi} \int \left( \Delta g - H \frac{d\Delta g}{d\rho} + \frac{H^2}{2} \frac{d^2 \Delta g}{d\rho^2} \right) s(\psi) d\psi + \frac{dT}{d\rho} H - \frac{1}{2} \frac{d^2 T}{d\rho^2} H^2 = T + \delta_3,$$

where

$$\delta_3 = \frac{H^3}{\rho} \left( \frac{1}{2} \frac{d\Delta g}{d\rho} - \frac{2\Delta g}{\rho} - \frac{6T}{\rho^2} \right).$$

It is easily understood that the quantity  $\delta_3$  containing  $H^3$  will not appear in the final result, if in formula (6) with  $i = 2$  we set  $\rho = a$ .

In conclusion, let us note that the process of calculations according to formula (4), as well as Molodenskiy's process of successive approximations, leads to accurate values of the disturbing potential on the sphere of radius  $a$  if only in the last of the used formulae (6) we assume  $\rho = a$ .

# Bibliography

1. Marych, M.I.: Ob opredelenii otkloneniy otvesa na fizicheskoy poverkhnosti Zemli. (On the Determination of Plumb Line Deflections on the Earth's Physical Surface). From symposium "Geodeziya, Kartografiya i aerofotos"yemka," No. 3, Lvov University, publishers, 1965.
2. Molodenskiy, M.S., Yeremeyev, V.F., Yurkina, M.I.: Metody izucheniya vneshnego gravitatsionnogo polya i figury zemli. (Methods of Investigation of the External Gravitational Field and Shape of the Earth). Trudy TsNIIGAIK, No. 131. Moscow, Geodezizdat, 1960.